INDEX

(New genera, species, etc., are printed in *italics*)

abdominalis, Hippocampus, 529.
Acanthobothrium, 134.
Acanthocephala, 124, 151.
acanthodes, Makrokylindrus, 435.
accessor, Castor, 285, 288.
Achaetonura euchaetiae, 382, 411.
  melalophae, 382, 411.
Achras, 356.
  calcicola, 357.
  calcicolafolia, 338, 340, 356, 359 (fig.).
  chicle, 357.
aciculatus, Perilampus, 370, 380.
Acmeodon, 229.
acmeodontoides, *Emperodon*, 223, 229.
Acoetidae, 265.
acolytus, Ellipsodon, 242.
Acraea andromacha, 252.
Acrepidopterum minutum, 197.
  minutum *apicalis*, 197.
Acroporidae, 90, 100.
Actinacis, 90, 92, 100.
  alabamiensis, 101.
  barretti, 73, 101, 110.
  sawkinsi, 73, 100, 110.
acuñai, Cyrtinus, 206.
adareanum, Nymphon, 418.
Adelometra tenuipes, 247.
Adisophanes miscellus, 50.
adkinsi, Synastrea, 73, 83, 87, 109.
admirabilis, Ptilodus, 225.
Admontia setigera, 18, 22.
Adocia, 444, 445.
  cinerea, 445, 455.
Adocus bossi, 181, 186.
  kirtlandius, 186.
advenus, Operculinoides, 487, 489, 495.
affinis, Paradidyma, 20, 21, 27, 35.
Agamonema *immanis*, 125, 142, 149.
  vomitor, 125, 150.
Agariciidae, 78, 96.
agaricites, Synastrea, 88.
agassizi, Leptophyllia, 73.
agnesae, Hippocampus, 529.
alabamensis, Dichocoenia, 75.
alabamiensis, Actinacis, 101.
alascana, *Retiometra*, 248.
alaskensis, Ammothea, 414, 421.
alba, Inga, 349.
albertensis, Boremys, 170–173.
albiguttus, Paralichthys, 148.
Alcyonium aurantium, 462.
aldrichi, Paradidyma, 21, 30.
aldrichi, Spanipalpus, 52.
alexinus, Perilampus, 370, 410, 411.
Alilepus, 111, 120.
  annectens, 119.
  vagus, 119 (fig.), 121, 285, 288.
Allamanda, 357.
alope, Cercyonis alope, 256.
alticuspis, Palaechthon, 223.
amblopiitis, Proteocephalus, 137–139.
Amblypoda, Fort Union, 224, 244.
Ambrosia, 405.
americana, Ficus, 341, 343.
  Rileya, 483.
americanafolia, Ficus, 341, 342 (fig.).
americanus, Mastodon, 286.
Amia calva, 137.
Ammothea, 414, 418.
  alaskensis, 414, 421.
  discoidea, 414, 417 (fig.), 418.
  latifrons, 420, 421.
Ammotheidae, 414.
Ammothella, 414, 421.
  longicaudata, 414, 421.
Amorpha fruticosa, 407.
Amoy, China, polychaetous annelids from, 261.
amoyensis, Nereis (Neanthes), 261, 269 (fig.), 272.
ampelus, Ernestia, 382, 411.
Amphibia, fossil, from Hagerman, Idaho, 285.
Amphicaecum *parvum*, 125, 143.
Amphicyon sp., 288.
Amphinomidae, 261.
Anabacia, 86.
Anabaciidae, 84.
Anacardiaceae, fossil Venezuelan, 352.
analis, Clinocottus, 326.
Anaptomorphidae, 223.
Anchoviella epsetus, 125, 144.
anchoviellae, Rhaphidascaris, 124, 125, 144.
Ancylis comptana, 401.
andina, Cuphocera, 46, 49, 63.
  Epicuphocera, 46, 63.
Andira, 349.
androcardia, Enodia, 252.
Andrognathus, 364.
  corticarius, 364.
andromacha, Acraea, 252.
  Enodia, 252.
  Enodia portlandia, 253, 254.
  Hipparchia, 252.
  Papilio (Oreas Marmorata), 252.
  Papilio (Parnassius), 252.

Anepsyra jaumei, 193.
Angiospermophyta, fossil Venezuelan, 344.
angulosa, Mycale, 448.
angusticornis, Atrophopalpus, 10, 15.
Ceratomyiella, 11, 15.
angustifolia, Sphaeralcia, 407.
angustus, Sympherobius, 395, 411.
Anisakinae, 124, 141.
Anisonchus sectorius, 224, 244.
Anisota senatoria, 378.
Annametra, 247.
minuta, 247.
occidentalis, 247.
annandalei, Clymene (Euclymene), 277.
Eyclymene, 277.
annectens, Allepus, 119.
Lepus, 119, 120.
Annelids, polychaetous, from Amoy, China, 261.
polychaetous, new species of Nereidae from California, 467.
anomalos, Faviosseris, 72, 82, 110.
anomocerus, Perilampus, 370, 371, 375, 398.
Anona, 345.
guppyi, 339, 340, 345.
macgravii, 347.
montana, 347.
sphaerocarpa, 347.
sphaerocarpoides, 338, 340, 346 (fig.), 347.
Anonaceae, fossil Venezuelan, 345.
Anonales, 345.
Anseriformes, fossil, from Idaho, 285.
antarctica, Eometra, 248.
Psathyrometra, 248.
Antedon clio, 248.
hageni(1), 245, 246.
petasus, 247.
Antedonidae, 245.
Antedoninae, 247.
Antennarius, 512.
anthedon, Enodia portlandia, 255, 256.
Anthocephalus macrourus, 131.
Antholithus venezuelensis, 338, 340.
antiguensis, Operculinoides, 487, 488, 492, 496.
antillarum, Antilloseris, 98.
Elaphidion, 192.
Antillia, 96.
Antillophyllia, 80, 96.
sp., 73, 96.
Antilloseris, 81, 96.
antillarum, 98.
cantabrigiensis, 73, 96.
cyclolites, 98.
jamaicaensis, 73, 97.
sp., 73, 97, 110.
antiqua, Gasterocoma, 6.
antiquorum, Hippocampus, 517, 518, 525, 530, 540, 551, 568, 570.
antiquus, Hippocampus, 518, 521, 525, 570.
apachensis, Hypolagus, 117.
Apanteles hyphantriae, 382, 411.
melanoscelus, 382, 411.
sp., 403, 411.
aperta, Paradidyma, 18, 21, 29.
Aphrodita flava, 261.
Aphronurus, 230.
fraudator, 223, 230.
apicalis, Acrepidopterum minutum, 197.
Paradidyma, 20, 21, 33.
Aplophyllia, 76.
Aplysilla, 443.
glacialis, 442, 443.
lendenfeldi, 443.
Apocynaceae, fossil Venezuelan, 357.
Apocynophyllum, 357.
salvadorensis, 338, 340, 357, 360.
texensis, 341.
apua, Mycteroperca, 154.
aquilonius, Ellipsodon, 224, 242.
Arachnida, pycnogonids from Puget Sound, 413.
Archaeolagus ennisianus, 117.
macrocephalus, 117.
primigenius, 117.
Archohelia, 105.
Arctocyonidae, 223, 232.
arcuata, Kathleena, 143.
Arecaceae, fossil Venezuelan, 344.
Arecales, 344.
arenarum, Camelops, 285, 288.
areolata, Nectandra, 340, 354.
Arhythmorhynchus, 153, 154.
duocinctus, 125, 152.
frassoni, 153.
fuscus, 153, 154.
hispidus, 153.
siluricola, 153.
aristalis, Paradidyma, 20, 28.
armata, Lachnommopsis, 18, 41.
Paradidyma, 20, 21, 41.
Arnoglossus, 134.
Artiodactyla, fossil, from Blanco, Texas, 288.
fossil, from Hagerman, Idaho, 285, 288.
Ascaridae, 141.
Ascaris sp., 148.
Ascogaster sp., 403, 411.
aspera, Sabicea, 360.
asperifolia, Sabicea, 340, 360.
asperrima, Florometra, 250.
Asphondylia opuntiae, 483.
Aspideretes austerus, 185.
fontanus, 185.
ovatus, 182 (fig.), 183, 186.
vegetus, 183.
vorax, 183, 184 (fig.), 185, 186.
Assilina, 489.
Asterosmilia, 106.
hilli, 104, 106.
Astraea decactis, 104.
Astraraea, 88.
media, 88.
Astreopora, 102.
walli, 73, 102, 110.

Astrocoenia, 95, 104.
decaturensis, 95.
duerdeni, 73.
jamaicaensis, 73, 95, 110.
Astrocoeniidae, 95.
Astrocottus, 330.
leprops, 330, 331 (fig.).
Atactorhynchus, 151.
verecundus, 123, 125, 151.
Atlantic Ocean, sponges from near Panama Canal, 455.
atrichus, Hippocampus, 524, 525, 540.
Atrophopalpus, 10.
angusticornis, 10, 15.
Atrophopoda, 17.
braueri, 9, 18.
peruana, 43.
singularis, 17, 18, 38.
townsendi, 10, 12.
atrosanguinea, Microciona, 442, 448.
aurantia, Tethya, 462.
aurantium, Alcyonium, 462.
aurea, Chiloepalpus (Cuphocera), 57, 70.
Cuphocera, 70.
aurifrons, Cuphocera, 63.
auritus, Phalacrocorax, 285.
austerus, Aspideretes, 185.
australis, Cuphocera, 48, 57.
australis, Proteocephalus, 125, 135, 139.
australis, Spanipalpus, 57.
Aves, fossil, from Hagerman, Idaho, 285.
azteca, Felis concolor, 214.
Baena, 169, 170, 174, 177, 187.
fluviatilis, 177.
hatcheri, 172.
nodosa, 166, 168, 169, 186.
ornata, 165, 166 (fig.), 167 (fig.), 186.
sp., 186.
Baenidae, 165, 186.
Bagre marina, 125, 130, 135, 146.
bairdii, Lepus bairdii, 112.
bakeri, Perilampus, 386, 388.
Barbados, fossil corals from, 103.
barbara, Spongia, 455.
barbinervis, Bignonia, 359.
barretti, Actinacis, 73, 101, 110.
Basilemys, 178, 181, 187.
nobilis, 178 (fig.), 179 (fig.), 180, 186.
praeciara, 180, 181.
sinuosa, 180, 181.
variolosa, 180, 181.
Bat, little brown, new trematode from, 321.
bathmodon, Pantolambda, 244.
Bathycuma longicaudata, 423.
Bathymetrinae, 248.
baueri, Neurankylus, 165, 186.
beameri, Cuphocera, 49, 66.
Beetles, new Cerambycidae from West Indies, 189.
belli, Chasmosaurus, 164.
Marphysa, 266.
beringi, Lamprops, 431.
Berry, Edward Wilbur, on Tertiary plants from Venezuela, 335.
betijoquensis, Blechnum, 338, 340.
Ficus, 340.
biaculeatus, Gastrotokeus, 515.
Syngnathoides, 515.
Bibliography (Literature Cited), on West Indian fossil corals, 107.
on Tertiary Foraminifera, 494.
on parasites of Galveston Bay fishes, 155.
on Chinese Polychaeta, 279.
on Puget Sound Pycnogonida, 422.
on Reptilia of Kirtland formation, 187.
on Panama sponges, 465.
on tapeworms from carnivores, 220.
on trematodes of subfamily Pleurogenetinae, 324.
on fossil vertebrates from Hagerman, Idaho, 315.
bicalcaratus, Platygonus, 288.
bicincta, Ceratomyiella, 11, 13.
bicolor, Loxogenes, 323.
bifascipennis, Callimome, 481.
Bigelovia, 407.
Bignonia, 358.
barbinervis, 359.
cujabamba, 359.
eximia, 359.
triphylla, 358.
zuliana, 338, 340, 358, 359 (fig.).
Bignoniaceae, fossil Venezuelan, 358.
biustus, Leptostylus, 204.
Blanco, Texas, fauna, 288.
Blarina gidleyi, 285.
Blechnum betijoquensis, 338, 340.
bleekeri, Hippocampus, 529.
Bloch, Marc Elieser, his account of seahorses, 514.
Boleometra, 248.
clio, 248.
Bolin, Rolf Ling, on new cottid fishes from western Pacific and revision of genus Stlengis, 325.
boltonae, Diplaraea, 73, 83, 109.
borealis, Enodia portlandia, 255, 256.
Boremys, 169, 170, 173, 187.
albertensis, 170–173.
grandis, 170, 171 (fig.), 172 (fig.), 173, 186.
pulchra, 169–173.
Borophagus, 287.
diversidens, 288.
sp., 285, 288.
bossi, Adocus, 181, 186.
boweni, Rhizophora, 338, 340.
Brachycybe, 361, 364.
lecontei, 363 (fig.), 365, 366.
lecontii, 366.
petasata, 363 (fig.), 365.
producta, 363 (fig.), 365, 367.
rosea, 363 (fig.), 365, 367, 368.
Brachylagus idahoensis, 112, 114–116.
Brachyphyllia, 89.
branchiatus, Cirratulus, 261, 274 (fig.), 276.

Brandesia, 323.
brasiliana, Paradidyma, 19, 38.
braueri, Atrophopoda, 9, 18.
Brenthis *hana*, 257.
breviceps, Hippocampus, 530.
brevirostris, Hippocampus, 499, 510, 520–522, 525, 546, 570, 571.
briareus, Taxocrinus, 4, 5.
browni, Hypolagus, 116, 117.
brunneofasciatus, Leptostylus, 205.
brunneus, Hippocampus, 568, 569.
buccata, Cuphocera, 46, 48, 59.
Bucephalidae, 126.
Bucephalopsis haimeana, 127.
Bucephalus papillosus, 127.
Burserites fayettensis, 341.
venezuelana, 338, 340.
Butterflies, genus Enodia and new fritillary from Peru, 251.
caballus, Equus, 292, 294, 301–303, 305–309, 311–314.
Cactus insects, new Chalcidoidea parasitic on, 481.
Caesalpinia, 349.
Caesalpiniaceae, fossil Venezuelan, 349.
Calamophyllia, 76.
calcicola, Achras, 357.
calcicolafolia, Achras, 338, 340, 356, 359 (fig.).
California, Cumacea from, 423.
new annelids (Nereidae) from, 467.
californica, Chrysopa, 395.
californica, Diastylis, 431, 432 (fig.).
Hemilamprops, 424, 429, 430 (fig.), 439.
californicus, Lynx rufus, 214, 217, 219.
Platydesmus, 367.
californiensis, Cuphocera, 70.
Callimome, 481.
bifasciipennis, 481.
Callimomidae, 481.
Calliobothrium, 134.
filicolle, 134.
calva, Amia, 137.
cambridgensis, Stylophora, 73, 95, 110.
Camelops arenarum, 285, 288.
campana, Hircinia, 456.
Spongia, 457.
Camptisocale, 460.
Campylaspis, 427.
canaliculata, 426 (fig.), 427, 438.
canadensis, Lynx, 219.
Perilampus, 370, 371, 374, 392, 411.
canaliculata, Campylaspis, 426 (fig.), 427, 438.
Canimartes cumminsii, 288.
Canis lestes, 217.
cantabrigiensis, Antilloseris, 73, 96.
Dendracis, 73, 100.
Turbinoseris, 96.
capitatus, Perilampus, 371, 375, 397.
Carabocrinus, 6.
carbonarium, Haemulon, 148.
Carcharias littoralis, 134.
cardui, Vanessa, 252.
Caribbean Sea, sponges from near Panama Canal, 441.
caribboea, Cliona, 461.
carinata, Cyclaspis, 427.
Lamprops, 431.
carinifrons, Perilampus, 370, 373, 388.
Carnivora, fossil, from Blanco, Texas, 288.
fossil, from Fort Union of Montana, 223, 232.
fossil, from Hagerman, Idaho, 285, 288.
new tapeworms from, 211.
carolinensis, Perilampus, 370, 373, 376, 410.
Carpolestidae, 223.
Caryometra, 247.
tenuipes, 247.
Caryophylliidae, 74.
Cassia, 349.
excelsa, 349.
longifolia, 340, 349.
spectabilis, 349.
zuliana, 340, 346 (fig.), 350.
Castor, 283.
accessor, 285, 288.
Castoroides, 286.
catadupensis, Mesomorpha, 72.
Trochoseris, 72, 78.
catadupensis, *Vaughanoseris*, 72, 80, 81, 109, 110.
Catemophrys sequens, 39.
caviractus, Pantolambda, 224, 244.
Ceanothus, 382.
cecilia, Mycale, 447, 448 (fig.).
Cedrola jacksoniana, 341, 342 (fig.).
celata, Cliona, 461.
Centrastrea, 80.
hilli, 72, 80, 109.
microphyllia, 80.
Centrorhynchidae, 152.
Centrorhynchus, 154.
Centrosaurus, 164.
centrura, Dasybatis, 133.
cepedianum, Dorosoma, 125, 143.
cephalus, Mugil, 125, 142.
Cerambycidae, new West Indian, 189.
Ceratomeryx prenticei, 285, 288.
Ceratomyiella angusticornis, 11, 15.
bicincta, 11, 13.
conica, 10, 11, 13–17.
orbitalis, 11, 16.
review of genus, 9, 10.
townsendi, 11, 12.
Ceratops, 160.
Ceratopsia, 160.
Ceratopsidae, 163, 185.
Cercyonis alope alope, 256.
alope nephele, 256.
alope pegala, 256.
Cervus, 283.
Cestoda, 129.
new species from carnivores, 211.
Chalcid flies of genus Perilampus, 369.
Chalcidoidea, 371.
new, parasitic on cactus insects, 481.
Chalcis violacea, 371.

Chandler, Asa Crawford, on parasites of fishes in Galveston Bay, Tex., 123.
Chasmosaurus belli, 164.
kaiseni, 186.
sp., 164 (fig.), 185.
Chelonia, 165.
Chen pressa, 285.
chicle, Achras, 357.
Chiloepalpus, 47, 70.
(Cuphocera) aurea, 57, 70.
China, polychaetous annelids from Amoy, 261.
Chisternon, 171.
undatum, 172.
Chloeia, 261.
flava, 261.
chloropus, Gallinula, 285.
Chondrilla, 464.
nucula, 464.
Chriacus, 233, 234, 236.
pelvidens, 234, 235.
pugnax, 223, 235.
pusillus, 223, 234, 235.
christianiaensis, Goniaraea, 73, 103, 110.
Chrysobalanus, 347.
icaco, 348.
preicaco, 348.
venezuelanus, 338, 340, 342 (fig.), 347.
Chrysopa californica, 395.
sp., 395, 411.
chrysopae, Perilampus, 371, 374, 394, 410, 411.
Chrysophyllum preoliviforme, 341.
Chusquea, 345.
cinerea, Adocia, 445, 455.
Reniera, 445.
Spongia, 444, 445.
cinerescens, Paradidyma, 20, 26.
Cirratulidae, 276.
Cirratulus, 276.
branchiatus, 261, 274 (fig.), 276.
Citharexylon retiforme, 347.
Cladocera jamaicaensis, 72.
Cladophyllia, 76.
Claenodon, 233.
ferox, 223, 232.
latidens, 223.
montanensis, 223.
silberlingi, 223, 232.
vecordensis, 223, 232.
claremontus, Gosodesmus, 363 (fig.), 364.
clarendonensis, Eupsammia, 73, 98, 99, 110.
Clark, Austin Hobart, on butterflies of genus Enodia and new fritillary from Peru, 251.
on new genera and species of unstalked crinoids, 245.
clinactina, Goniaraea, 103.
Clinocottus analis, 326.
clio, Antedon, 248.
Boleometra, 248.
Cliona, 461.
caribboea, 461.
celata, 461.
Clymene, 277.
(Euclymene) annandalei, 277.
Coccometra hagenii, 245–247.
cochranensis, Ectypodus, 226.
cockburnensis, Nereis, 469.
coerulescens, Haliclona, 444, 445.
Reniera, 444.
Coleoptera, new West Indian Cerambycidae, 189.
Cole, W. Storrs. (See under Vaughan, T. W.)
collaris, Lysidice, 261, 266.
collieri, Contracaecum, 124, 125, 141, 142.
Colobognatha, new species of, 361.
columbia, Trypespongia, 456.
columnaris, Dictuophyllia conferticostata, 72.
Columnastrea eyrei, 73.
Colurostylis, 439.
occidentalis, 423, 431, 439.
Colymbiformes, fossil, from Idaho, 285.
Colymbus sp., 285.
Combretaceae, fossil Venezuelan, 355.
Combretum, 355.
incertum, 356.
petraflumensis, 356.
stephensoni, 338, 340, 355, 359 (fig.).
Cominia occidentalis, 247.
comma, Conoryctes, 223.
complanata, Operculina, 489.
Compsilura concinnata, 382, 411.
Compsometra, 245, 246.
nuttingi, 245, 246.
parviflora, 247.
comptana, Ancylis, 401.
concinnata, Compsilura, 382, 411.
concordiarcensis, Prodiacodon, 223, 228.
Condaminea, 360.
grandifolia, 340, 360.
Condylarthra, Fort Union, 224, 238.
conferticosta, Leptoria, 77.
conferticostata, Dictuophyllia, 72, 77.
Diploria, 77.
Leptoria, 77.
conformis, Cuphocera, 48, 54.
conica, Ceratomyiella, 10, 11, 13–17.
Conoryctes comma, 223.
conradi, Eupsammia, 100.
contigua, Cuphocera, 46, 49, 61, 66.
contorta, Stylophora, 73, 94.
Contracaecum, 142.
collieri, 124, 125, 141, 142.
quadricuspe, 143.
robustum, 125, 142.
microcephalum, 142, 143.
micropapillatum, 141, 143.
spiculigerum, 143.
tricuspe, 143.
cooki, Cremastus, 403, 411.
cookii, Lutravus, 285, 288.
Copecrypta, 47.
nitens, 70.
nitidifrons, 70.
cora, Crotalocrinus, 6.
Corals, fossil, from West Indies, 71.
cordatus, Piperites, 340, 345, 346 (fig.).
Cordillerion tropicus, 288.

coriacea, Persea, 340, 355.
Coriphagus montanus, 224.
corticarius, Androgynathus, 364.
Corynecrinus, new Devonian crinoid genus, 1, 4.
romingeri, 1, 2, 5.
costata, Cyclaspis, 427.
Cottidae, from western Pacific, 325.
Cottontail, sage (Sylvilagus nuttalli grangeri), 112, 114.
cotylophora, Dichelyne, 147.
Coussapoa villosoides, 340.
crassicollis, Taenia, 217, 218.
crassiseta, Paradidyma, 20, 27.
crassolamellosa, Diploria, 77.
Crateroseris, 82.
crawfordi, Perilampus, 370, 373, 384.
Cremastus, 398, 411.
cooki, 403, 411.
creola, Debis, 253, 254.
Enodia, 253–257.
Cretaceous, Upper, corals of Jamaica, 72, 74.
Crinoids, fossil, new genus of, 1.
unstalked, new genera and species of, 245.
Crocodilia, 187.
Crotalocrinus cora, 6.
Crustacea, California Cumacea, 423.
Cryptotrema, 323.
Cucullanellus, subg., 149.
Cucullanidae, 146.
Cucullanus lintoni, 148.
cujabamba, Bignonia, 359.
Cumacea, California, 423.
cumminsii, Canimartes, 288.
Equus, 289.
Plesippus, 288.
cunctatrix, Spirastrella, 461.
cunea, Hyphantria, 382.
Cuphocera andina, 46, 49, 63.
aurea, 70.
aurifrons, 63.
australis, 48, 57.
beameri, 49, 66.
buccata, 46, 48, 59.
californiensis, 70.
conformis, 48, 54.
contigua, 46, 49, 61, 66.
erythrostoma, 70.
flavicornis, 49, 58.
fucata, 46, 49, 63, 65.
geminata, 48, 55.
hirsuta, 46, 48, 49, 54, 62, 63, 65, 66.
incongrua, 46, 48, 68.
macrocera, 47, 50, 52, 70.
miscelli, 48, 49, 52.
nitidifrons, 70.
parksi, 48, 50, 52, 59.
revision of American species of, 45.
ruficauda, 70.
scutellaris, 48, 53.
stricklandi, 63.
torosa, 46, 48, 67.
cuspidata, *Eudaemonema*, 223, 231.
Cuvier, Georges, his account of seahorses, 518, 521.
cyaneus, Perilampus, 370, 410.
cyathiformis, Heliastraea, 90.
Multicolumnastraea, 73, 90.
Cyathocrinidae, 4.
Cyathocrinoidea, 1, 4–6.
Cyathocrinus, 5.
glaber, 5.
Cyathomorpha, 89.
Cyathoseris, 72.
haidingeri, 72.
Cycadaceae, fossil Venezuelan, 344.
Cycadales, 344.
Cycadophyta, fossil Venezuelan, 344.
Cyclaspis, 423, 424.
carinata, 427.
costata, 427.
levis, 425, 427.
longipes, 427.
nubila, 424 (fig.).
picta, 427.
pusilla, 427.
sibogae, 427.
unicornis, 427.
varians, 427.
Cyclolites, 84, 86.
discoidea, 85.
hemisphaerica, 85.
jamaicaensis, 72, 84, 109.
ligeriensis, 85.
michelini, 85.
cyclolites, Antilloseris, 98.
Cycloseris, 85, 86.
Cygnus sp., 285.
cylindricus, Dichelyne, 148.
Cynipsillum, 371.
Cynodontomys, 231.
Cynoscion regalis, 134, 144.
Cyprinodon, 124, 142.
variegatus, 123–125, 140–142, 150, 152.
cyprinodontis, *Glossocercus*, 123, 125, 140.
Cyrtinus *acuñai*, 206.
eugeniae, 207.
Cysticercoides *menidiae*, 125, 141.
Dasybatis centrura, 133.
Dasychone, 278.
orientalis, 278.
Datana integerrima, 378.
dawsoni, Diastylopsis, 423, 436, 437.
Debis creola, 253, 254.
portlandia, 253.
decactis, Astraea, 104.
Madracis, 103, 104, 110.
decaturensis, Astrocoenia, 95.
decipiens, Glycera, 275.
Deinodontidae, 160, 186.
delauneyi, Nanilla, 197.
deliculata, Leptoria, 78.
Deltatheridiidae, 223, 227.
Dendracis, 100.
cantabrigiensis, 73, 100.
dentatus, Paralichthys, 148.
Deopalpus, 46.
hirsutus, 46, 63.
depressidens, Paromomys, 223.
derelicta, Paradidyma, 19, 25, 26, 28.

Dermatemydidae, 165, 178, 186.
Dermosmilia, 84.
deserticola, Lepus californicus, 112.
Deuterogonodon, 232.
montanus, 223, 232, 233.
Devonian, Clark County, Ind., new crinoid genus from, 1.
dia, Operculinoides, 493, 494.
Diaphoropeza, 18.
peruana, 31.
Diastylis, 431.
californica, 431, 432 (fig.).
Diastylopsis dawsoni, 423, 436, 437.
tenuis, 436, 437 (fig.).
diazi, Romerolagus, 120.
Dichelyne, 149.
cotylophora, 147.
cylindricus, 148.
diplocaecum, 125, 149.
fastigatus, 125, 146.
fossor, 147.
mauritanicus, 148.
robusta, 147.
Dichocoenia, 75.
alabamensis, 75.
stokesi, 75.
trechmanni, 72, 75, 109.
tuberosa, 75.
Dicotyledonae, 345.
Dictuophyllia, 77.
conferticostata, 72, 77.
conferticostata columnaris, 72.
Didelphodontinae, 227.
Didelphodus, 227.
Didyma validinervis, 17, 37.
Didymictis, 237, 238.
haydenianus, 224, 237.
microlestes, 224, 238.
tenuis, 224, 238.
Dilepididae, 139.
Dilepis kempi, 141.
diluculi, Palaeosinopa, 223, 230.
Dimorpharaea, 82.
Dimorphostylis, 435.
Dinosauria, 160, 185.
Diopatra, 266.
neapolitana, 266.
Dioscorea, 345.
Diplaraea, 83.
boltonae, 73, 83, 109.
venezuelensis, 84.
Diploastrea, 89.
diplocaecum, Dichelyne, 125, 149.
diploderma, Tethya, 451.
Diplolepis violacea, 371.
Diploria conferticostata, 77.
crassolamellosa, 77.
Diprion, 382.
Diptera, review of flies of genera Ceratomyiella and Paradidyma, 9.
revision of American Cuphocera, 45.
hosts of Perilampus, 411.
disceptatrix, *Haplaletes*, 224, 243, 244.
discoidea, Ammothea, 414, 417 (fig.), 418.
discoidea, Cyclolites, 85.
disjunctus, *Litaletes*, 224, 242.
dissentaneus, *Litomylus*, 224, 243.
distoechus, Stiengis, 325, 327, 328–330, (fig.).
Distomum monticelli, 128.
sp., 128.
distorta, Nereis (Leptonereis), 261, 273, 274 (fig.).
diversidens, Borophagus, 288.
Dolichohippus, 294.
dolomieu, Micropterus, 137.
doria, Haliclona, 458.
Dorosoma cepedianum, 125, 143.
Douglass, Earl, work in Montana, 221.
douglassi, Ptilodus, 223, 225.
Dryopteris, 343.
duerdeni, Astrocoenia, 73.
duocinctus, Arhythmorhynchus, 125, 152.
durissima, Strongylophora, 459.
eakini, Nereis (Nereis), 468, 472, 473 (fig.), 476.
Ebenales, 356.
Echeneibothrium, 134.
Echinorhynchus medius, 154.
Ectocion, 240, 241.
Ectypodus, 226.
cochranensis, 226.
grangeri, 223, 226.
musculus, 226.
russelli, 223, 226.
silberlingi, 223, 225, 226.
ecuadorensis, Hippocampus, 534, 537.
edensis, Hypolagus, 116.
Edentata, fossil, from Blanco, Texas, 288.
fossil, from Hagerman, Idaho, 285, 288.
egle, Euchaetias, 382.
eifelianus, Lecythocrinus, 4–6.
Elachipalpus macrocera, 52.
Elaphidion antillarum, 192.
lanatum, 192, 193.
manni, 193.
monticola, 191.
wickhami, 192.
elegans, Heliconia, 340.
elisa, *Plakoosa*, 463 (fig.).
elizabethae, *Paracycloseris*, 73, 86, 109, 110.
Ellipsodon, 241, 242, 243.
acolytus, 242.
aquilonius, 224, 242.
elongatus, Proteocephalus, 125, 137.
Sciocyrtinus, 208.
Elphidotarsius florencae, 223.
Emperodon, 229.
acmeodontoides, 223, 229.
ennisianus, Archaeolagus, 117.
Enodia androcardia, 252.
andromacha, 252.
creola, 253–257.
notes on butterflies of genus, 251.
portlandia, 253, 254, 256, 257.
portlandia andromacha, 253, 254.
portlandia *anthedon*, 255, 256.
portlandia *borealis*, 255, 256.
portlandia portlandia, 255, 256.

Entada, 335, 341.
entadaformis, Leguminosites, 338, 340.
entellus, Perilampus, 370, 410, 411.
Eocene, corals of Jamaica, 72, 93, 94.
Venezuelan plants, 335, 341.
Eometra, 248.
antarctica, 248.
Epalpus, 70.
Epiblema strenuana, 405.
Epicuphocera, 46.
andina, 46, 63.
epsetus, Anchoviella, 125, 144.
Equus, 281, 283, 286, 289, 290, 293, 295, 298, 299, 310–313.
caballus, 292, 294, 301–303, 305–309, 311–314.
cumminsii, 289.
excelsus, 283, 292.
grevyi, 294, 295, 300–303, 305–309, 311–314.
idahoensis, 283, 291, 292.
pacificus, 291.
scotti, 282, 316–320.
erectus, Hippocampus, 516, 525, 561, 566.
Syngnathus, 516, 517.
erina, Haliclona, 457.
erinaceus, Tetrarhynchus, 132.
Ernestia ampelus, 382, 411.
ruficauda, 382, 411.
Erythrina, 349.
erythrostoma, Cuphocera, 70.
Essostrutha ramsdeni, 210.
roberto, 209.
eucapitis, Nereis (Nereis), 468, 469 (fig.).
euchaetiae, Achaetonura, 382, 411.
Euchaetias egle, 382.
Euclymene, 277.
annandalei, 277.
Eucosmodon sp., 223.
Eudaemonema, 231.
cuspidata, 223, 231.
Eugenia, 341.
eugeniae, Cyrtinus, 207.
Eulimneria valida, 382, 411.
sp., 382, 411.
Eulophidae, 485.
Eulophus, 407.
Eupatorium serotinum, 395.
Eupogonius *haitiensis*, 197, 199.
pilosulus, 198.
wickhami, 199.
Euprotogonia, 240.
minor, 239.
puercensis, 239.
Eupsammia, 99.
clarendonensis, 73, 98, 99, 110.
conradi, 100.
Eupsammidae, 99.
europaeus, Hippocampus, 499, 510, 511, 528, 530, 531, 541–543, 546, 547 (fig.), 571, 572.
Europe, seahorses of, 497.
Eurytomidae, 482.
Eusmiliidae, 75.
excelsa, Cassia, 349.
excelsus, Equus, 283, 292.
exigua, Retiometra, 248.
eximia, Bignonia, 359.
Exline, Harriet I., on pycnogonids from Puget Sound, 413.
exsculpta, Heliastraea, 90.
eyrei, Columnastrea, 73.
Fagara, 350.
fagifolia, Inga, 349.
falconeri, Rhynchotherium, 288.
Farlapiscis, 530.
Farr, Marcus Stults, work in Montana, 221.
fasciatus, Lynx, 217, 219.
Lynx fasciatus, 217, 219.
Lynx rufus, 214.
fascicularis, Hippocampus, 561, 567.
fastigatus, Dichelyne, 125, 146.
fastuosa, Inga, 349.
Faviidae, 76, 96.
Favioseris, 82.
anomalos, 72, 82, 110.
fayettensis, Bursérites, 341.
Felis concolor azteca, 214.
concolor hippolestes, 212, 214.
concolor oregonensis, 214.
hillanus, 288.
lacustris, 285, 288.
lynx, 217, 219.
felis, Galeichthys, 125, 130, 131, 133, 135, 154.
femoratum, Nymphon, 421.
Phoxichilidium, 414, 421.
fenestratus, Pentaceratops, 160, 163, 185.
ferox, Claenodon, 223, 232.
ferrea, *Fisherispongia*, 460 (fig.).
Ficus, 357.
americana, 341, 343.
americanafolia, 341, 342 (fig.).
betijoquensis, 340.
jynx, 343.
laqueata, 342.
mississippiensis, 354.
pseudomediafolia, 342.
wilcoxensis, 343.
filicolle, Calliobothrium, 134.
Synbothrium, 130.
Syndesmobothrium, 131.
Fisher, Warren Samuel, on new West Indian cerambycid beetles, 189.
Fisherispongia, 460.
ferrea, 460 (fig.).
Fishes, new cottid, and revision of genus Stlengis, 325.
Galveston Bay, parasites of, 123.
review of seahorses (Hippocampus) of American and European coasts, 497.
flagelliformis, Inga, 349.
flava, Aphrodita, 261.
Chloeia, 261.
flavicornis, Cuphocera, 49, 58.
flexuossissima, Leptoria, 77.
Flies, American, of genus Cuphocera, 45.
muscoid, review of genera Paradidyma and Ceratomyiella, 9.

florencae, Elphidotarsius, 223.
floridensis, Nummulites, 489.
Florometra asperrima, 250.
fluviatilis, Baena, 177.
Thescelus, 177.
foliatus, Hippocampus, 515.
Syngnathus, 515, 516.
fontanus, Aspideretes, 185.
Foraminifera, Tertiary, of genera Operculina and Operculinoides from North America and West Indies, 487.
forresti, Operculinoides, 488, 493, 495.
Fort Union of Montana, new Paleocene mammals from, 221.
Fossils, corals from West Indies, 71.
new Devonian crinoid, 1.
new Tertiary Foraminifera from North America and West Indies, 487.
hares from late Pliocene of southern Idaho, 111.
horse remains from upper Pliocene of Idaho, 281.
new mammals from Fort Union (Paleocene) of Montana, 221.
Tertiary Venezuelan plants, 335.
Reptilia of Kirtland formation of New Mexico, 159.
fossor, Dichelyne, 147.
fragile, Synbothrium, 130.
Syndesmobothrium, 130.
francescana, Pliohippus, 289.
frassoni, Arhythmorhynchus, 153.
fraudator, Aphronurus, 223, 230.
Fritillary, new species from Peru, 251.
fruticosa, Amorpha, 407.
fucata, Cuphocera, 46, 49, 63, 65.
Trichophora, 65.
fulvicornis, Perilampus, 370, 371, 376, 402, 411.
Fundulus, 124.
heteroclitus, 125, 140, 142, 148, 150.
Fungia, 86.
(Cycloseris) patella, 110.
furcatus, Ictalurus, 125, 149, 150.
furens, Prothryptacodon, 223, 233, 234.
furlongi, Hypolagus, 118 (fig.), 121.
fusca, Sabella, 278.
Sabellastarte, 278.
fuscus, Arhythmorhynchus, 153, 154.
Gahan, Arthur Burton, on new Chalcidoidea parasitic on cactus insects, 481.
gahani, Perilampus, 371, 375, 401.
Galeichthys felis, 125, 130, 131, 133, 135, 154.
milberti, 133, 134.
Gallinula chloropus, 285.
Galveston Bay, Tex., parasites of fishes from, 123.
Gasterocoma, 6.
antiqua, 6.
Gasterocomidae, 4–6.
Gasterostomum gracilescens, 127.
sp., 126.
Gastrotokeus biaculeatus, 515.
gavialidis, Goezia, 146.
Gazin, Charles Lewis, on fossil hares from late Pliocene of southern Idaho, 111.
on fossil horse remains from upper Pliocene of Idaho, 281.
Gelastops, 227.
parcus, 223, 227.
geminata, Cuphocera, 48, 55.
Gentialales, 357.
Geodia, 454.
gibberosa, 454, 455.
Geraniales, 351.
Gerstaeckeria porosa, 486.
gerstaeckeriae, Tetrastichus, 485.
gibber, Gorgorhynchus, 125, 154.
gibberosa, Geodia, 454, 455.
Gidley, James Williams, work in Montana, 221.
Gidleya, 240.
gidleyi, Blarina, 285.
gidleyi, Ptilodus, 223, 225.
gidleyi, Thomomys, 285, 288.
Gidleyina, 240.
montanensis, 224, 240, 241.
silberlingi, 224, 240.
gigas, Gymnorhynchus, 125, 130, 133.
Gilmore, Charles Whitney, on Reptilia of Kirtland formation of New Mexico, with descriptions of new species of fossil turtles, 159.
Ginsburg, Isaac, on review of seahorses (Hippocampus), 497.
glaber, Cyathocrinus, 5.
glacialis, Aplysilla, 442, 443.
Simplicella, 443.
Glossocercus, 124, 139.
cyprinodontis, 123, 125, 140.
Glycera, 275.
decipiens, 275.
rouxii, 275.
Glyceridae, 275.
Glyptoporus, 321, 323.
noctophilus, 321, 322 (fig.).
Glyptotherium texanum, 288.
Goezia gavialidis, 146.
minuta, 125, 146.
goldmani, Ochotona schisticeps, 112.
Goniaraea, 103.
christianiaensis, 73, 103, 110.
clinactina, 103.
Goniopora, 90, 110.
jamaica, *I*, 90.
reussiana, 73, 90, 110.
trechmanni, 73, 91, 109.
Goniopteris, 343.
Gorgorhynchus, 154.
gibber, 125, 154.
medius, 154.
Gorgosaurus, 186.
libratus, 160.
sp., 160, 186.
Gosodesmus, 364.
claremontus, 363 (fig.), 364.
gracile, Nymphon, 418.
gracilescens, Gasterostomum, 127.
gracilis, Hippocampus, 534, 537.
Ptilodus, 225.

grandifolia, Condaminea, 340, 360.
grandis, Boremys, 170, 171 (fig.), 172 (fig.), 173, 186.
grandis, Spalacopsis, 201.
grangeri, Ectypodus, 223, 226.
grangeri, Sylvilagus nuttalli, 111, 114–116.
granulosus, Perilampus, 371, 375, 399.
grevyi, Equus, 294, 295, 300–303, 305–309, 311–314.
griseus, Neomaenis, 148.
Gruiformes, fossil, from Idaho, 285.
Gryporhynchus, 140.
guppyi, Anona, 339, 340, 345.
guttulatus, Hippocampus, 499, 500, 510, 514, 518, 521, 523, 525, 536, 537, 540–544, 548, 549, 571, 572.
Hippocampus guttulatus, 528, 530, 531, 543.
Gymnocanthus, 326.
Gymnorhynchus, 131.
gigas, 125, 130, 133.
malleus, 125, 130–132.
platycephalus, 133.
reptans, 131.
Hadrosauridae, 160, 161, 185.
Hadrosaurinae, 161.
haemorrhoa, Palpibraca, 46.
Haemulon, 148.
carbonarium, 148.
hageni(i), Antedon, 245, 246.
Coccometra, 245–247.
Hagerman, Idaho, fossil fauna, 288.
haidingeri, Cyathoseris, 72.
haimeana, Bucephalopsis, 127.
haitiensis, Eupogonius, 197, 199.
Halichondria, 449, 457.
panicea, 442, 449, 455.
Haliclona, 444, 445, 457.
coerulescens, 444, 455.
doria, 458.
erina, 457.
permollis, 442, 444, 445.
halieus, Pelecanus, 285.
Halina, 462.
Halinidae, 462.
Halisarca lobularis, 455.
hana, Brenthis, 257.
Haplaletes, 243.
disceptatrix, 224, 243, 244.
Haplaraea, 84.
Haplomylus, 242, 243.
hardyi, Nereis (Eunereis), 480.
Hares, fossil, from late Pliocene of southern Idaho, 111.
Hartman, Olga, on new polychaetous annelids of family Nereidae from California, 467.
Haruspex inscriptus, 195, 196.
insularis, 195, 196.
similis, 196.
hatcheri, Baena, 172.
haydenianus, Didymictis, 224, 237.
Heliastraea cyathiformis, 90.
exsculpta, 90.
Heliconia elegans, 340.
Hemilamprops, 424, 429.
californica, 424, 429, 430 (fig.), 439.
uniplicata, 431.
hemionus, Odocoileus, 214.
Odocoileus hemionus, 214.
hemisphaerica, Cyclolites, 85.
hemispherica, Thescelus, 174, 175 (fig.), 176 (fig.), 177, 186.
Hemiuridae, 127.
hemuloni, Synbothrium, 130.
henekeni, Paracyathus, 105.
heptagonus, Hippocampus, 515, 525, 570.
Hernandia, 354.
tongi, 340, 353 (fig.), 354.
Hernandiaceae, fossil Venezuelan, 354.
hesperus, Machairodus, 285, 288.
heteracanthum, Pterobothrium, 131.
heteroclitus, Fundulus, 125, 140, 142, 148, 150.
heteropoda, Lumbriconereis, 266.
Lumbrinereis, 266.
hiemale, Nymphon, 416.
hildebrandi, Hippocampus, 511, 529, 536, 578, 579, 580 (fig.), 581 (fig.).
hillanus, Felis, 288.
hilli, Asterosmilia, 104, 106.
hilli, Centrastrea, 72, 80, 109.
hilli, Trochosmilia, 78, 79.
Hipparchia andromacha, 252.
Hippocampus abdominalis, 529.
agnesae, 529.
antiquorum, 517, 518, 525, 530, 540, 551, 568, 570.
antiquus, 518, 521, 525, 570.
atrichus, 524, 525, 540.
bleekeri, 529.
breviceps, 530.
brevirostris, 499, 510, 520–522, 525, 546, 570, 571.
brunneus, 568, 569.
ecuadorensis, 534, 537.
erectus, 516, 525, 561, 566.
europaeus, 499, 510, 511, 528, 530, 531, 541–543, 546, 547 (fig.), 571, 572.
fascicularis, 561, 567.
foliatus, 515.
gracilis, 534, 537.
guttulatus, 499, 500, 510, 514, 518, 521, 523, 525, 536, 537, 540–544, 548, 549, 571, 572.
guttulatus guttulatus, 528, 530, 531, 543.
guttulatus *multiannularis*, 499, 501, 523–525, 528, 530, 531, 536-538 (fig.), 539 (fig.), 540, 544, 545, 548–550.
heptagonus, 515, 525, 570.
hildebrandi, 511, 529, 536, 578, 579, 580 (fig.), 581 (fig.).
hudsonius, 499, 500, 510, 512, 518, 527, 536, 537, 544, 545, 548, 549, 551, 564, 566, 578, 583.
hudsonius hudsonius, 528, 530, 533, 550 (fig.), 551, 552 (fig.), 553 (fig.), 554 (fig.), 569.

Hippocampus hudsonius kincaidi, 528, 530, 533, 544, 557, 566, 568, 574.
ingens, 506, 527, 529–531, 534, 535 (fig.), 549, 579–582.
jubatus, 524, 525, 570.
kincaidi, 501, 511, 568.
laevicaudatus, 551, 560.
longirostris, 519–523, 525, 543, 546, 572.
marginalis, 561, 567.
non aculeatus, incisuris raris, 543.
obtusus, 511, 529, 576, 577 (fig.), 579, 581, 582.
pentagonus, 530.
poeyi, 561, 567.
punctulatus, 499, 500, 517, 518, 525, 527, 551, 561, 568, 572, 575, 582, 583, 593.
ramulosus, 511, 517, 518, 525, 543, 546.
regulus, 506, 529, 558, 584, 586 (fig.), 587 (fig.), 590.
reidi, 528, 530, 532, 536, 537, 564, 566, 569, 571–573 (fig.), 574 (fig.).
review of genus, 497.
rondeletii, 519.
rosaceus, 521, 525, 543.
rosamondae, 589, 593, 594.
stylifer, 499, 561, 566, 567.
trimaculatus, 517, 518.
villosus, 529, 582, 583.
zosterae, 502, 506, 512, 525, 529, 585, 588, 589.
Hippocampus, subg., 527, 529, 530.
hippocampus, Syngnathus, 513–520, 525, 530, 543, 570.
hippolestes, Felis concolor, 212, 214.
Hircinia, 456.
campana, 456.
variabilis, 457.
hirsuta, Cuphocera, 46, 48, 49, 54, 62, 63, 65, 66.
hirsutus, Deopalpus, 46, 63.
hispidus, Arhythmorhynchus, 153.
Hooks, method of measurement of in cestodes, 211.
Horses, fossil, from upper Pliocene of Idaho, 281.
hudsonius, Hippocampus, 499, 500, 510, 512, 518, 527, 536, 537, 544, 545, 548, 549, 551, 564, 566, 578, 583.
Hippocampus hudsonius, 528, 530, 533, 550 (fig.), 551, 552 (fig.), 553 (fig.), 554 (fig.), 569.
Hyaenognathus, 287.
sp., 285, 288.
hyalinus, Perilampus, 369, 370, 380, 410, 411.
Hymenoptera, chalcid flies of genus Perilampus, 369.
hosts of Perilampus, 411.
Hyopsodontidae, 224, 241.
Hyopsodus, 242.
simplex, 242.
Hypacrosaurus, 162.
Hyphantria cunea, 382.
hyphantriae, Apanteles, 382, 411.
Meteorus, 382, 411.
Hypolagus, 111, 114, 116, 119, 120.
apachensis, 117.
browni, 116, 117.
edensis, 116.
furlongi, 118 (fig.), 121.
limnetus, 113 (fig.), 114, 115 (fig.), 118, 121, 285, 288.
vetus, 112 (fig.), 114, 116, 117, 119–121, 285, 288.
Hypsioma *insularis*, 199.
picticornis, 200.
icaco, Chrysobalanus, 348.
Icelus, 326.
Ictalurus furcatus, 125, 149, 150.
Ictidopappus, 237.
mustelinus, 224, 237.
Idaho, fossil hares from late Pliocene, 111.
fossil horse remains from upper Pliocene of, 281.
idahoensis, Brachylagus, 112, 114–116.
Equus, 283, 291, 292.
Ischyrosmilus, 283.
Lutravus, 285, 288.
Phalacrocorax, 285.
Pseudemys, 285.
ignis, Tedania, 459.
Thalysias, 459.
immanis, Agamonema, 125, 142, 149.
imperfecta, Mycale, 448.
incertum, Combretum, 356.
incongrua, Cuphocera, 46, 48, 68.
Indrodon, 231.
Inga, 348.
alba, 349.
fagifolia, 349.
fastuosa, 349.
flagelliformis, 349.
pinetorum, 349.
reissi, 340, 346 (fig.), 348.
sp., 338, 340, 349.
tetraphylla, 349.
ingens, Hippocampus, 506, 527, 529–531, 534, 535 (fig.), 549, 579–582.
inscriptus, Haruspex, 195, 196.
Insectivora, fossil, Fort Union, 223, 227.
fossil, from Hagerman, Idaho, 285.
Insects, butterflies of genus Enodia and new fritillary from Peru, 251.
chalcid flies of genus Perilampus, 369.
new Chalcidoidea parasitic on cactus insects, 481.
new West Indian cerambycid beetles, 189.
insilens, Thescelus, 174–177.
insularis, Hypsioma, 199.
Haruspex, 195, 196.
integerrima, Datana, 378.
intermedia, Placospongia, 454, 455.
intermedius, Pantolambda, 224, 244.
inversus, Pentacodon, 230.

iortlandia, Vanessa, 252.
Ischyrosmilus idahoensis, 283.
Isodictya permollis, 444.
itea, Vanessa, 252.
jacksoniana, Cedrola, 341, 342 (fig.).
Jamaica, fossil corals from, 72, 74, 93, 94.
jamaica *1*, Goniopora, 90.
jamaicaensis, Antilloseris, 73, 97.
jamaicaensis, Astrocoenia, 73, 95, 110.
jamaicaensis, Cladocera, 72.
jamaicaensis, Cyclolites, 73, 84, 109.
jamaicaensis, Stiboriopsis, 72.
Turbinoseris, 97.
Jamsus, subg., 525, 529, 571, 583, 584.
japonicus, Thelepus, 278.
jasminifolium, Pleonotoma, 358.
jaumei, Anepsyra, 193.
jepseni, Parectypodus, 223, 227.
jubatus, Hippocampus, 524, 525, 570.
jynx, Ficus, 343.
kaiseni, Chasmosaurus, 186.
Kathleena arcuata, 143.
kempi, Dilepis, 141.
kincaidi, Hippocampus, 501, 511, 568.
Hippocampus hudsonius, 528, 530, 533, 544, 557, 566, 568, 574.
Kirk, Edwin, on Corynecrinus, a new Devonian crinoid genus, 1.
Kirtland fauna and its geological age, 185.
Kirtland formation of New Mexico, Reptilia from, 159.
kirtlandius, Adocus, 186.
koninckl, Leptoria, 78.
Kritosaurus, 160, 162, 186, 187.
navajovius, 161, 185.
Lachnomma, 18.
magnicornis, 9, 18, 38, 39.
Lachnommopsis, 18.
armata, 18, 41.
lacustris, Felis, 285, 288.
ladae, Leptacodon, 223, 228.
laevicaudatus, Hippocampus, 551, 560.
laevicephalus, Perilampus chrysopae, 394, 395.
laevis, Perilampus, 370, 411.
Lagomorpha, fossil, from Hagerman, Idaho, 285, 288.
Lambeosaurinae, 161.
Lamprops, 424, 431.
beringi, 431.
carinata, 431.
lanatum, Elaphidion, 192, 193.
lanceolata, Styrax, 340, 356.
Tapirira, 340, 352, 353 (fig.), 360.
laqueata, Ficus, 342.
Lastrea, 343.
laticollis, Taenia, redescription of, 211, 216.
latidens, Claenodon, 223.
latidens, Tricentes, 224, 236.
latifrons, Ammothea, 420, 421.
latrunculus, *Spanoxyodon*, 224, 236.
Laubenfels, Max Walker de, on Panama sponges, 441.
Lauraceae, fossil Venezuelan, 354.
Laurales, 354.
Laxosuberites, 450.
zeteki, 450, 451 (fig.), 455.
Leach, William Elford, his account of seahorses, 517.
Lecithochirium *microstomum*, 125, 127.
synodi, 128.
lecontei, Brachycybe, 363 (fig.), 365, 366.
lecontii, Brachycybe, 366.
Lecythocrinidae, fossil crinoid family, 1, 4.
Lecythocrinus, 2, 4–6.
eifelianus, 4–6.
Leguminosae, fossil Venezuelan, 351.
Leguminosites entadaformis, 338, 340.
venezuelensis, 338, 340.
Leiostomus xanthurus, 125, 148, 154.
Leipsanolestes, subg., 228.
lemhi, Ochotona princeps, 112.
lendenfeldi, Aplysilla, 443.
Leodicidae, 266.
lepida, Tentacularia, 125, 129.
Lepidasthenia, 264.
ocellata, 261, 263 (fig.), 264.
Lepidium, 407.
Lepidocyclina macdonaldi, 490.
mantelli, 494.
tobleri, 490.
trinitatis, 490.
Lepidonotus, 262.
minutus, 261, 262, 263 (fig.).
Lepisosteus, 124, 185.
osseus, 125, 137, 139.
Leporidae, fossil, from southern Idaho, 111.
lepreos, Perilampus, 370, 411.
leprops, *Astrocottus*, 330, 331 (fig.).
Leptacodon, 228.
ladae, 223, 228.
munusculum, 223, 228.
packi, 228.
siegfriedti, 228.
tener, 228.
Leptictidae, 223, 228.
Leptonereis, subg., 275.
leptonyx, Megalonyx, 285, 288.
Leptophyllia, 84.
agassizi, 73.
Leptophylliidae, 83.
Leptoria conferticosta, 77.
conferticostata, 77.
deliculata, 78.
flexuossissima, 77.
koninckl, 78.
reticulata, 78.
leptostomus, Megalonyx, 288.
Leptostylus biustus, 204.
brunneofasciatus, 205.
maculifer, 205.
monticola, 201.
ornatus, 203.
planicollis, 203.
pygmaeus, 206.
vanduzeei, 204.
lepturus, Trichiurus, 125, 128, 145.

Lepus, 114, 116, 117, 120, 286.
annectens, 119, 120.
bairdii bairdii, 112.
californicus deserticola, 112.
townsendii, 113, 120.
townsendii townsendii, 112.
lestes, Canis, 217.
lethostigmus, Paralichthys, 124, 125, 142, 153.
levis, Cyclaspis, 425, 427.
libratus, Gorgosaurus, 160.
ligeriensis, Cyclolites, 85.
lignea, Spongia, 455.
Limatulum, 323.
limitare, Siphonophora, 362, 363 (fig.).
limnetus, Hypolagus, 113 (fig.), 114, 115 (fig.), 118, 121, 285, 288.
lindgreni, Neotragoceras, 283.
linea, Nereis (Neanthes), 261, 268, 269 (fig.).
Linnaeus, Carolus, his account of seahorses, 513.
lintoni, Cucullanus, 148.
Lintoniella, 130.
Lissodendoryx, 446, 447.
Litaletes, 242–244.
disjunctus, 224, 242.
Litharaea, 91, 92.
Litomylus, 243.
dissentaneus, 224, 243.
littoralis, Carcharias, 134.
Littorina sp., 124.
lobularis, Halisarca, 455.
Oscarella, 442, 455.
longicaudata, Ammothella, 414, 421.
Bathycuma, 423.
longifolia, Cassia, 340, 349.
longipes, Cyclaspis, 427.
longipes, Nereis (Eunereis), 467, 479, 480 (fig.).
longirostris, Hippocampus, 519–523, 525, 543, 546, 572.
Syngnathus, 522.
Loomis, Harold Frederick, on millipeds of order Colobognatha, 361.
Lophopsetta maculata, 148.
Loxogenes, 323.
bicolor, 323.
Lower California, millipeds from, 361.
lucifugus, Myotis, 321, 323.
Lumbriconereis heteropoda, 266.
Lumbrinereis, 266.
heteropoda, 266.
Lutra (Satherium) piscinaria, 285, 288.
Lutravus, 287.
cookii, 285, 288.
idahoensis, 285, 288.
lyncis, Taenia, 211, 212.
Lynx canadensis, 219.
fasciatus, 217, 219.
fasciatus fasciatus, 217, 219.
lynx, 217, 219.
rufus, 215, 217, 219.
rufus californicus, 214, 217, 219.
rufus fasciatus, 214.
rufus rufus, 214.
rufus uinta, 214.
lynx, Felis, 217, 219.
Lynx, 217, 219.
Lysidice, 266.
collaris, 261, 266.
lyttoni, Siphonacme, 364.
macdonaldi, Lepidocyclina, 490.
macgravii, Anona, 347.
Machairodus hesperus, 285, 288.
Macleayina, subg., 506, 526, 527, 536.
Macremphytus, 393.
Macrocentrus pallisteri, 403, 411.
sp., 405, 411.
macrocephalus, Archaeolagus, 117.
macrocera, Cuphocera, 47, 50, 52, 70.
Elachipalpus, 52.
Tachina, 52.
macrocystis, Taenia, 214, 217, 219.
macrourus, Anthocephalus, 131.
Odocoileus virginianus, 214.
maculata, Lophopsetta, 148.
maculifer, Leptostylus, 205.
Macy, Ralph W., on new trematode from little brown bat, 321.
Madracis, 104.
decactis, 103, 104, 110.
sp., 104, 105, 110.
magnicornis, Lachnomma, 9, 18, 38, 39.
Makrokylindrus, 435.
acanthodes, 435.
malacosomae, Tetrastichus, 485, 486.
Maldanidae, 277.
malleum, Synbothrium, 133.
malleus, Gymnorhynchus, 125, 130–132.
Mammals, fossil, from Hagerman, Idaho, 285.
fossil hares from late Pliocene of southern Idaho, 111.
new Paleocene from Fort Union of Montana, 221.
manni, Elaphidion, 193.
mantelli, Lepidocyclina, 494.
marana, Sophora, 338, 340, 346 (fig.), 350.
marginalis, Hippocampus, 561, 567.
mariannensis, Operculina, 489.
marina, Bagre, 125, 130, 135, 146.
marinus, Tylosurus, 126.
marmorata, Oreas, 252.
Marphysa, 266.
belli, 266.
orientalis, 261, 263 (fig.), 266.
sinensis, 261, 266.
Mastodon, 283.
americanus, 286.
mirificus, 283.
Matley, Charles Alfred, his collections of fossil corals, 74, 94.
matleyi, Trochocyathus, 72, 74, 109.
matthewi, Pronothodectes, 223.
maturus, Paromomys, 223.
mauritanicus, Dichelyne, 148.
media, Astraraea, 88.
mediator, Nereis, 469, 471.
medius, Echinorhynchus, 154.
Gorgorhynchus, 154.
Megalonyx leptostomus, 288.
leptonyx, 285, 288.

Megopterna minuta, 224.
melalophae, Achaetonura, 382, 411.
melanoscelus, Apanteles, 382, 411.
Melittia, 384.
Menidia, 127.
  menidia, 125, 126, 128, 141, 144.
  notata, 128.
menidia, Menidia, 125, 126, 128, 141, 144.
menidiae, Cysticercoides, 125, 141.
Meniscium, 343.
  palustre, 343.
  reticulatum, 343.
  wolfi, 340, 342 (fig.), 343.
meridionale, Nymphon, 416.
Mesomorpha catadupensis, 72.
Metachriacus, 235.
  provocator, 224, 235.
  punitor, 223, 235.
Meteorus hyphantriae, 382, 411.
  sp., 411.
mexicana, Microchira, 18, 29.
Miacidae, 224, 237.
michelini, Cyclolites, 85.
microcephalum, Contracaecum, 142, 143.
Microchira, 18.
  mexicana, 18, 29.
Microciona, 448.
  atrosanguinea, 442, 448.
microlestes, Didymictis, 224, 238.
Micropalpus, 70.
  ruficornis, 45, 46.
micropapillatum, Contracaecum, 141, 143.
microphyllia, Centrastrea, 80.
Micropogon undulatus, 125, 131, 154.
Microperus dolomieu, 137.
Microsmilia, 81.
Microsolena, 86.
microstomum, Lecithochirium, 125, 127.
milberti, Galeichthys, 133, 134.
Millipeds, new species of order Colobognatha, 361.
Mimomys, 287.
  primus, 285, 288.
Mimosaceae, fossil Venezuelan, 348.
miniata, Patiria, 474.
minor, Euprotogonia, 239.
  Ondatra idahoensis, 285, 288.
  Palenochtha, 223, 231.
minuta, *Annametra*, 247.
minuta, Goezia, 125, 146.
minuta, Megopterna, 224.
minutum, Acrepidopterum, 197.
minutus, Lepidonotus, 261, 262, 263 (fig.).
Miocene, Venezuelan plants, 336, 343.
miocenica, Pleonotoma, 338, 340, 358, 359 (fig.).
miocenica, Simaruba, 338, 340.
Mioclaenus, 241.
mirificus, Mastodon, 283.
misakia, Schmidtia, 325, 330.
  Schmidtina, 330.
  Stlengis, 327, 330.
miscelli, Cuphocera, 48, 49, 52.
  Spanipalpus, 49.
  Trichophora, 46, 49.
miscellus, Adisophanes, 50.
mississippiensis, Ficus, 354.
Mixodectes, 231.
Mixodectidae, 223, 231.
Monellema ulkei, 485.
monellemae, Neocatolaccus, 484.
Monoclonius, 160.
Monocotyledonae, 344.
Montana, new Paleocene mammals from Fort Union of, 221.
montana, Anona, 347.
montanensis, Claenodon, 223.
montanensis, *Gidleyina*, 224, 240, 241.
montanensis, Myrmecoboides, 223.
montanus, Coriphagus, 224.
montanus, *Deuterogonodon*, 223, 232, 233.
montanus, Ptilodus, 223, 225.
monticellii, Distomum, 128.
monticola, Elaphidion, 191.
  Leptostylus, 201.
Mosesia, 323.
muesebecki, Perilampus, 371, 376, 407.
Mugil cephalus, 125, 142.
multiannularis, Hippocampus guttulatus, 499, 501, 523–525, 528, 530, 531, 536–538 (fig.), 539 (fig.), 540, 544, 545, 548–550.
Multicolumnastraea, 90.
  cyathiformis, 73, 90.
  parvula, 90.
multifragum, Psittacotherium, 223.
Multituberculata, Fort Union, 223, 224.
munusculum, Leptacodon, 223, 228.
Muscidae, review of flies of genera Ceratomyiella and Paradidyma, 9.
musculus, Ectypodus, 226.
mustelinus, *Ictidopappus*, 224, 237.
Mycale, 447.
  angulosa, 448.
  cecilia, 447, 448 (fig.).
  imperfecta, 448.
  phyllophila, 448.
Mycteroperca apua, 154.
Myledaphus, 185.
Mylodon sp., 288.
Myotis lucifugus, 321, 323.
Myoxocephalus, 326.
Myriapoda, new millipeds of order Colobognatha, 361.
Myrmecoboides montanensis, 223.
Myrtales, 355.
Myxilla, 446.
Nanilla delauneyi, 197.
  tuberculata, 196.
Nannippus phlegon, 288.
natans, Nereis (Nereis), 468, 474, 475 (fig.).
navajovius, Kritosaurus, 161, 185.
neapolitana, Diopatra, 266.
Nectandra, 354.
  areolata, 340, 354.
Nematoda, 141.

Neocatolaccus, 484.
monellemae, 484.
tylodermae, 484.
Neoechinorhynchidae, 151.
Neohipparion sp., 288.
Neomaenis, 148.
griseus, 148.
neomexicana, Paradidyma, 19, 21, 23.
neonigripes, Nereis (Nereis), 468, 471, 472 (fig.), 474, 476.
Neotragoceras lindgreni, 283.
nephele, Cercyonis alope, 256.
Nephthydidae, 276.
Nephthys, 276.
sinensis, 261, 276.
Nereidae, 268.
new species from California, 467.
Nereis, 268, 467, 468.
(Neanthes) *amoyensis*, 261, 269 (fig.), 272.
cockburnensis, 469.
(Leptonereis) *distorta*, 261, 273, 274 (fig.).
(Nereis) *eakini*, 468, 472, 473 (fig.), 476.
(Nereis) *eucapitis*, 468, 469 (fig.).
(Eunereis) hardyi, 480.
(Neanthes) *linea*, 261, 268, 269 (fig.).
(Eunereis) *longipes*, 467, 479, 480 (fig.).
mediator, 469, 471.
(Nereis) *natans*, 468, 474, 475 (fig.).
(Nereis) *neonigripes*, 468, 471, 472 (fig.), 474, 476.
(Neanthes) *orientalis*, 261, 269 (fig.), 270.
(Alitta) oxypoda, 268.
(Neanthes) oxypoda, 261, 268.
pelagica, 467, 472.
procera, 468, 471.
(Nereis) *pseudoneanthes*, 470 (fig.).
(Neanthes) *saltoni*, 477, 478 (fig.).
(Ceratonereis) *tunicatae*, 476, 477 (fig.).
vexillosa, 469–471.
(Neanthes) virens, 479.
Neurankylus, 187.
baueri, 165, 186.
Neuroptera, hosts of Perilampus, 411.
New Mexico, Reptilia from Kirtland formation of, 159.
nitens, Copecrypta, 70.
nitidifrons, Copecrypta, 70.
Cuphocera, 70.
nitidus, Perilampus canadensis, 371, 374, 393.
nobilis, Basilemys, 178 (fig.), 179 (fig.), 180, 186.
noctophilus, *Glyptoporus*, 321, 322 (fig.).
nodosa, Baena, 166, 168, 169, 186.
North America, chalcid flies of genus Perilampus north of Mexico, 369.
new Foraminifera from Tertiary of, 487.
North and South America, seahorses of, 497.
notata, Menidia, 128.
nubila, Cyclaspis, 424 (fig.).
nubriterrae, Peromyscus maniculatus, 214.
nucula, Chondrilla, 464.
Nummilites floridensis, 489.
sp., 490.
willcoxi, 489.
nutrix, Rhabdophylliа, 77.
nuttingi, Compsometra, 245, 246.
Nyctitheriidae, 223, 229.
Nymphon, 414.
adareanum, 418.
femoratum, 421.
gracile, 418.
hiemale, 416.
meridionale, 416.
solitarium, 414, 417 (fig.).
tridentatum, 418.
turritum, 414, 416, 417 (fig.).
Nymphonidae, 413.
obliqua, Paradidyma, 20, 22.
obtusus, Hippocampus, 511, 529, 576, 577 (fig.), 579, 581, 582.
occidentalis, *Annametra*, 247.
Colurostylis, 423, 431, 439.
Cominia, 247.
ocellata, Lepidasthenia, 261, 263 (fig.), 264.
ocellatus, Perilampus, 370, 373, 390.
ocellatus, Sciaenops, 124, 125, 147, 148.
Ochotona princeps lemhi, 112.
schisticeps goldmani, 112.
octonemus, Polynemus, 154.
Oculina, 105.
Odocoileus columbianus scaphiatus, 214.
hemionus, 214.
hemionus hemionus, 214.
virginianus, 214.
virginianus macrourus, 214.
Oedemapeza, 10.
townsendi, 12.
Onchobothrium, 134.
Ondatra idahoensis minor, 285, 288.
operculella, (Phthorimaea) Gnorimoschema, 401.
Operculina complanata, 489.
mariannensis, 489.
new Tertiary North American and West Indian, 487, 488.
tuberculata, 487, 488, 495.
Operculinella, 489.
venosa, 489.
Operculinoides *advenus*, 487, 489, 495.
antiguensis, 487, 488, 492, 496.
dia, 493, 494.
forresti, 488, 493, 495.
new Tertiary North American and West Indian, 487, 489.

Operculinoides *semmesi*, 487, 488, 491, 495, 496.
tuxpanicus, 488, 494, 495.
vicksburgensis, 487, 490, 495.
Ophlitaspongiidae, 460.
Opuntia, 482.
opuntiae, Asphondylia, 483.
opuntiae, Rileya, 482.
Orasema viridis, 372.
orbitalis, Ceratomyiella, 11, 16.
Oreas marmorata, 252.
oregonensis, Felis concolor, 214.
orientalis, Dasychone, 278.
orientalis, Marphysa, 261, 263 (fig.), 266.
Nereis (Neanthes), 261, 269 (fig.), 270.
ornata, Baena, 165, 166 (fig.), 167 (fig.), 186.
Pseudoeme, 189.
ornatipennis, Spalacopsis, 200.
ornatus, Leptostylus, 203.
Orthonopias triacis, 326.
osborni, Pliohippus, 289.
Oscarella, 455.
lobularis, 442, 455.
osensis, Stlengis, 325–328.
osseus, Lepisosteus, 125, 137, 139.
Oulastrea, 89.
Oulastreidae, 88.
ovatus, Aspideretes, 182 (fig.), 183, 186.
oxypoda, Nereis (Alitta), 268.
Nereis (Neanthes), 261, 268.
Oxyurostylis, 423, 437.
pacifica, 429, 437, 438 (fig.).
smithi, 423, 437, 438.
Pacific Ocean, sponges from near Panama Canal, 443.
western, new cottid fishes from, 325.
pacifica, Oxyurostylis, 429, 437, 438 (fig.).
pacificus, Equus, 291.
packi, Leptacodon, 228.
Palaechthon, 231, 232.
alticuspis, 223.
Palaeolagus, 117.
Palaeosinopa, 230.
diluculi, 223, 230.
Palenochtha, 231.
minor, 223, 231.
Paleocene, new mammals from Fort Union of Montana, 221.
pallisteri, Macrocentrus, 403, 411.
Palmophyllum, 344.
sp., 340, 344.
Palpibraca, 46.
haemorrhoa, 46.
palustre, Meniscium, 343.
Panama Canal, sponges from, 441.
panamensis, Panthalis, 265.
Pavona, 106.
Pandosentis, 151.
panicea, Halichondria, 442, 449, 455.
Spongia, 449.
Panthalis, 265.
panamensis, 265.
Pantolambda, 244.
bathmodon, 244.
cavirictus, 224, 244.
intermedius, 224, 244.
Pantolambdidae, 224, 244.
Pantolestidae, 223, 230.
Papilio (Parnassius) andromacha, 252.
(Oreas Marmorata) andromacha, 252.
portlandia, 251.
Papilionaceae, fossil Venezuelan, 350.
papillosus, Bucephalus, 127.
Parabascus, 323.
Paracyathus, 104, 105.
henekeni, 105.
sp., 72, 74.
Paracycloseris, 85.
elizabethae, 73, 86, 109, 110.
Paradiastylis, 435.
Paradidyma *affinis*, 20, 21, 27, 35.
aldrichi, 21, 30.
aperta, 18, 21, 29.
apicalis, 20, 21, 33.
aristalis, 20, 28.
armata, 20, 21, 41.
brasiliana, 19, 38.
cinerescens, 20, 26.
crassiseta, 20, 27.
derelicta, 19, 25, 26, 28.
neomexicana, 19, 21, 23.
obliqua, 20, 22.
peruana, 21, 31, 38.
(Diaphoropeza) peruana, 43.
peruviana, 43.
petiolata, 19, 21, 39.
piliventris, 21, 32.
retracta, 19, 26.
review of genus, 9, 17.
setigera, 20, 22, 23.
singularis, 10, 15, 19, 21, 31, 32, 36–38, 41.
townsendi, 12.
validinervis, 20, 23, 25, 29, 33–35, 37, 39, 40, 42, 43.
Paralichthys albiguttus, 148.
dentatus, 148.
lethostigmus, 124, 125, 142, 153.
Parasaurolophus, 160, 162, 186.
tubicen, 161, 185.
walkeri, 162, 163.
Parasites, chalcid flies of genus Perilampus, 369.
Chalcidoidea on cactus insects, 481.
of fishes in Galveston Bay, Tex., 123.
Paraxiothea, 277.
parcus, *Gelastops*, 223, 227.
Parectypodus, 227.
jepseni, 223, 227.
simpsoni, 227.
tardus, 225.
parksi, Cuphocera, 48, 50, 52, 59.
Paromomys, 231, 232.
depressidens, 223.
maturus, 223.
parviflora, Compsometra, 247.

parvula, Multicolumastraea, 90.
parvum, Amphicaecum, 125, 143.
patella, Fungia (Cycloseris), 110.
Patiria miniata, 474.
Pavona, 106.
panamensis, 106.
pennyi, 106.
sp., 104, 106, 110.
pegala, Cercyonis alope, 256.
pelagica, Nereis, 467, 472.
Pelecaniformes, fossil, from Idaho, 285.
Pelecanus halieus, 285.
Peleteria, 47, 70.
pelvidens, Chriacus, 234, 235.
pennyi, Pavona, 106.
Pentaceratops, 160, 163, 186.
fenestratus, 160, 163, 185.
sternbergii, 160, 163, 185.
Pentacodon, 230.
inversus, 230.
pentacus, Protogonodon, 233.
pentagonus, Hippocampus, 530.
Perilampidae, 371.
Perilampus aciculatus, 370, 380.
alexinus, 370, 410, 411.
anomocerus, 370, 371, 375, 398.
bakeri, 386, 388.
canadensis, 370, 371, 374, 392, 411.
canadensis *nitidus*, 371, 374, 393.
capitatus, 371, 375, 397.
carinifrons, 370, 373, 388.
carolinensis, 370, 373, 376, 410.
chrysopae, 371, 374, 394, 410, 411.
chrysopae laevicephalus, 394, 395.
crawfordi, 370, 373, 384.
cyaneus, 370, 410.
entellus, 370, 410, 411.
fulvicornis, 370, 371, 376, 402, 411.
fulvicornis *prothoracicus*, 371, 376, 403.
gahani, 371, 375, 401.
granulosus, 371, 375, 399.
hyalinus, 369, 370, 380, 410, 411.
laevis, 370, 411.
lepreos, 370, 411.
muesebecki, 371, 376, 407.
ocellatus, 370, 373, 390.
platigaster, 383.
platygaster, 369, 370, 373, 383, 410.
regalis, 370, 372, 378.
revision of genus (in America north of Mexico), 369.
rohweri, 371, 375, 396.
robertsoni, 371, 376, 409.
similis, 371, 376, 406.
stygicus, 371, 376, 404, 411.
subcarinatus, 370, 373, 386.
triangularis, 411.
Periptychidae, 224.
Periptychus, 244.
Perissodactyla, fossil, from Blanco, Texas, 288.
fossil, from Hagerman, Idaho, 285, 288.
permollis, Haliclona, 442, 444, 445.
Isodictya, 444.
Peromyscus maniculatus nubriterrae, 214.
Perry, George, his account of seahorses, 515.
Persea, 355.
coriacea, 340, 355.
sp., 338, 340, 355.
Personales, 358.
Peru, new fritillary from, 251.
peruana, Atrophopoda, 43.
Diaphoropeza, 31.
Paradidyma, 21, 31, 38.
Paradidyma (Diaphoropeza), 43.
peruviana, Paradidyma, 43.
petasata, Brachycybe, 363 (fig.), 365.
petasus, Antedon, 247.
petiolata, Paradidyma, 19, 21, 39.
petraflumensis, Combretum, 356.
Phalacrocorax auritus, 285.
idahoensis, 285.
sp., 285.
Phaneropsolus, 323.
Phenacodontidae, 224, 238.
phlegon, Nannippus, 288.
Phorbasidae, 446.
Phoreiobothrium, 134.
triloculatum, 134.
Phoxichelidiidae, 414.
Phoxichilidium, 414, 421.
femoratum, 414, 421.
(Phthorimaea) Gnorimoschema operculella, 401.
Phyllobothrioidea, 134.
phyllophila, Mycale, 448.
Phyllopteryx, 516.
Physoseris, 84.
Phytoadmontia, 18.
setigera, 22.
Phytophaga sp., 482.
Picrodus silberlingi, 224.
picta, Cyclaspis, 427.
picticornis, Hypsioma, 200.
Pika (Ochotona princeps lemhi and O. schisticeps goldmani), 112.
piliventris, Paradidyma, 21, 32.
pilosulus, Eupogonius, 198.
pinetorum, Inga, 349.
Piper, 345.
Piperaceae, fossil Venezuelan, 345.
Piperales, 345.
Piperites, 345.
cordatus, 340, 345, 346 (fig.).
Pisces, fossil, 185, 187.
fossil, from Hagerman, Idaho, 285.
(See also under Fishes.)
piscinaria, Lutra (Satherium), 285, 288.
Pithecolobrium, 349.
Placospongia, 454.
intermedia, 454, 455.
Plakoosa, 462.
elisa, 463 (fig.).
Plakortis, 462.
planicollis, Leptostylus, 203.
Plants, Tertiary Venezuelan, 335.
Plastomenus robustus, 186.
sp., 186.
platigaster, Perilampus, 383.

platycephalus, Gymnorhynchus, 133.
Tetrarhynchus, 130, 131, 133.
platygaster, Perilampus, 369, 370, 373, 383, 410.
Platydesmus, 367.
californicus, 367.
Platygonus bicalcaratus, 288.
sp., 285.
texanus, 288.
Pleonotoma, 358.
jasminifolium, 358.
miocenica, 338, 340, 358, 359 (fig.).
tetraquetrum, 358.
Plesiadapidae, 223.
Plesippus, 281, 287–289.
cumminsii, 288.
proversus, 293, 294.
shoshonensis, 110, 111, 281, 284, 285, 288–314 (figs.), 316–320.
simplicidens, 282, 288, 289, 292, 293, 316, 318–320.
Plesippus quarry, near Hagerman, Idaho, fossil hares from, 114, 119.
Pleurogenes, 323.
Pleurogenetinae, key to, 323.
Pleurogenoides, 323.
pleuronectis, Scolex, 125, 134.
Pleurosternidae, 165, 186.
Pliauchenia spatula, 288.
pliciferus, Tetraclaenodon, 239.
Pliocene, late, fossil hares from southern Idaho, 111.
upper, of Idaho, fossil horse remains from, 281.
Pliohippus, 289, 290, 292.
francescana, 289.
osborni, 289.
proversus, 289.
Plumeria, 357.
Poacites, 344.
sp., 338, 340, 344.
Poales, 344.
Podoseris, 81.
poeyi, Hippocampus, 561, 567.
Polychaeta, new species of Nereidae from California, 467.
from Amoy, China, 261.
polymorphus, Scolex, 134.
Polynemus octonemus, 154.
Polynoidae, 262.
Polyphyllastrea, 82.
Polypodiaceae, fossil Venezuelan, 343.
Polypodiales, 343.
poolei, Pseudoeme, 190.
Populus, 382.
Porifera, Panama, 441.
Porites, 91.
reussiana, 90.
Poritidae, 90, 103.
porosa, Gerstaeckeria, 486.
Porrocaecum, 145.
secundum, 125, 145.
trichiuri, 125, 145.
portlandia, Debis, 253.
Enodia, 253, 254, 256, 257.
Enodia portlandia, 255, 256.
portlandia, Papilio, 251.
Vanessa, 252.
Postorchigenes, 323.
Poteriocrinus, 1.
praeclara, Basilemys, 180, 181.
praecursor, Serridentinus, 288.
preicaco, Chrysobalanus, 348.
prenticei, Ceratomeryx, 285, 288.
preoliviforme, Chrysophyllum, 341.
pressa, Chen, 285.
Prestonia, 357.
Primates, Fort Union, 231.
primigenius, Archaeolagus, 117.
primus, Mimomys, 285, 288.
Proboscidea, fossil, from Blanco, Texas, 288.
fossil, from Hagerman, Idaho, 285, 288.
Procamelus, 283, 285, 288.
Procampylaspis, 423, 429.
sp., 429.
procera, Nereis, 468, 471.
Prodiacodon, 228, 229.
concordiarcensis, 223, 228.
puercensis, 228.
Prodiploastrea, 88.
schindewolfi, 73, 88, 89, 110.
producta, Brachycybe, 363 (fig.), 365, 367.
Pronothodectes matthewi, 223.
Prosotocus, 323.
Prosthorhynchus, 154.
protelaria, Stenosmilia, 76.
Proteocephalidae, 135.
Proteocephalus, 124.
ambloplitis, 137–139.
australis, 125, 135, 139.
elongatus, 125, 137.
prothoracicus, Perilampus fulvicornis, 371, 376, 403.
Prothryptacodon, 233.
furens, 223, 233, 234.
Protogonodon, 232, 233.
pentacus, 233.
Protohippus, 283, 289.
Protoselene, 240, 241, 243.
proversus, Plesippus, 293, 294.
Pliohippus, 289.
provocator, *Metachriacus*, 224, 235.
proxima, Toxadocia, 442, 445.
Psathyrometra antarctica, 248.
Pseudemys idahoensis, 285.
Pseudoeme *ornata*, 189.
poolei, 190.
Pseudofavia, 89.
Pseudolmedia, 343.
pseudomediafolia, Ficus, 342.
pseudoneanthes, Nereis (Nereis), 470 (fig.).
Pseudosuberites, 449.
sulcatus, 442, 449.
Psittacotherium multifragum, 223.
Pteridophyta, fossil Venezuelan, 343.
Pterobothrium, 131.
heteracanthum, 131.
Pteromalidae, 484.

Ptilodontidae, 223, 224.
Ptilodus, 225.
admirabilis, 225.
douglassi, 223, 225.
gidleyi, 223, 225.
gracilis, 225.
montanus, 223, 225.
sinclairi, 223, 225, 226.
trovessartianus, 225.
puercensis, Euprotogonia, 239.
Prodiacodon, 228.
Puget Sound, pycnogonids from, 413.
pugnax, Chriacus, 223, 235.
pulchra, Boremys, 169–173.
punctulatus, Hippocampus, 499, 500, 517, 518, 525, 527, 551, 561, 568, 572, 575, 582, 583, 593.
punitor, *Metachriacus*, 223, 235.
pusilla, Cyclaspis, 427.
pusillus, Chriacus, 223, 234, 235.
Pycnogonids, from Puget Sound, 413.
pygmaeus, Leptostylus, 206.
Pylaie, ——— de la, his account of seahorses, 524.
quadricuspe, Contracaecum, 143.
quaylei, Rhabdophyllia, 72, 76, 109.
Querquedula sp., 285.
Rabbit, black-tailed jack (Lepus californicus deserticola), 112.
pigmy (Brachylagus idahoensis), 112, 114.
snow-shoe (Lepus bairdii bairdii), 112.
white-tailed jack (Lepus townsendii townsendii), 112.
Rafinesque, Constantine Samuel, his account of seahorses, 515.
ramsdeni, Essostrutha, 210.
ramulosus, Hippocampus, 511, 517, 518, 525, 543, 546.
rapiens, Thescelus, 174, 175, 177.
regalis, Cynoscion, 134, 144.
regalis, Perilampus, 370, 372, 378.
regulus, Hippocampus, 506, 529, 558, 584, 586 (fig.), 587 (fig.), 590.
reidi, Hippocampus, 528, 530, 532, 536, 537, 564, 566, 569, 571–573 (fig.), 574 (fig.).
Reinhard, Henry Jonathan, on revision of American Cuphocera, 45.
on American muscoid flies of genera Ceratomyiella and Paradidyma, 9.
reissi, Inga, 340, 346 (fig.), 348.
Reniera cinerea, 445.
coerulescens, 444.
reptans, Gymnorhynchus, 131.
Reptilia, fossil, from Hagerman, Idaho, 285.
fossil, from Kirtland formation of New Mexico, 159.
reticulata, Leptoria, 78.
reticulatum, Meniscium, 343.
retiforme, Citharexylon, 347.
Retiometra, 248.
alascana, 248.
exigua, 248.
retracta, Paradidyma, 19, 26.
reussiana, Goniopora, 73, 90, 110.
Porites, 90.
Rhabdophyllia, 76.
nutrix, 77.
quaylei, 72, 76, 109.
sp., 77.
Rhadinorhynchus tenuicornis, 123, 125, 154.
Rhaphidascaris, 144, 145.
anchoviellae, 124, 125, 144.
Rhinoceros, 283.
Rhipidocotyle *transversale*, 125, 126, 128, 141.
Rhizophora boweni, 338, 340.
Rhynchobothrium speciosum, 130.
Rhynchotherium falconeri, 288.
Richmond formation of Jamaica, lower Eocene, corals from, 93.
Ricuzenius, 331.
Rileya, 482.
americana, 483.
opuntiae, 482.
similaris, 482.
Risso, Antoine, his account of seahorses, 521.
roberto, Essostrutha, 209.
robertsoni, Perilampus, 371, 376, 409.
robusta, Dichelyne, 147.
robustum, Contracaecum, 125, 142.
robustus, Plastomenus, 186.
Rodentia, fossil, from Hagerman, Idaho, 285, 288.
rohweri, Perilampus, 371, 375, 396.
Romanes collection of fossil corals from Barbados, 103.
Romerolagus, 115, 120.
diazi, 120.
romingeri, *Corymecorinus*, 1, 2, 5.
rondeletii, Hippocampus, 519.
Roosa zyggompha, 462.
Rosaceae, fossil Venezuelan, 347.
rosaceus, Hippocampus, 521, 525, 543.
Rosales, 347.
rosamondae, Hippocampus, 589, 593, 594.
rosea, Brachycybe, 363 (fig.), 365, 367, 368.
rouxii, Glycera, 275.
Rubiaceae, fossil Venezuelan, 360.
Rubiales, 360.
ruficauda, Cuphocera, 70.
Ernestia, 382, 411.
ruficornis, Micropalpus, 45, 46.
rufus, Lynx, 215, 217, 219.
Lynx rufus, 214.
russelli, Ectypodus, 223, 226.
Sabella, 278.
fusca, 278.
Sabellastarte, 278.
fusca, 278.
Sabellidae, 278.
Sabicea, 360.
aspera, 360.
asperifolia, 340, 360.
saltoni, Nereis (Neanthes), 477, 478 (fig.).

salvadorana, Sophora, 338, 340.
salvadorensis, Apocynophyllum, 338, 340, 357, 360.
santa, Strongylophora, 459.
Sapindales, 352.
Sapotaceae, fossil Venezuelan, 356.
sawkinsi, Actinacis, 73, 100, 110.
scaphiatus, Odocoileus columbianus, 214.
schindewolfi, *Prodiploastrea*, 73, 88, 89, 110.
Schineria, 70.
Schinz, Heinrich Rudolf, his account of seahorses, 520.
Schmidtia, 325, 326.
misakia, 325, 330.
Schmidtina, 326.
misakia, 330.
schmiedeliana, Velates, 95, 101.
Sciaenops ocellatus, 124, 125, 147, 148.
Sciocyrtinus, 207.
elongatus, 208.
Scolex, 124.
pleuronectis, 125, 134.
polymorphus, 134.
Scotland beds of Barbados, corals from, 103.
scotti, Equus, 282, 316–320.
scutellaris, Cuphocera, 48, 53.
Sea-bean (Entada), 335, 341.
Seahorses, review of Hippocampus of American continents and Europe, 497.
sectorius, Anisonchus, 224, 244.
secundum, Porrocaecum, 125, 145.
Sematethmos, 72.
semmesi, Operculinoides, 487, 488, 491, 495, 496.
senatoria, Anisota, 378.
sequens, Catemophrys, 39.
Seriatoporidae, 94.
serotinum, Eupatorium, 395.
Serpentes, fossil, from Idaho, 285.
Serridentinus praecursor, 288.
setigera, Admontia, 18, 22.
Paradidyma, 20, 22, 23.
Phytoadmontia, 22.
shoshonensis, Plesippus, 110, 111, 281, 284, 285, 288–314 (figs.), 316–320.
sibogae, Cyclaspis, 427.
Siderastrea, 82.
Siderofungia, 82.
siegfriedti, Leptacodon, 228.
Silberling, Albert C., work in Montana, 221.
silberlingi, Claenodon, 223, 232.
silberlingi, Ectypodus, 223, 225, 226.
Gidleyina, 224, 240.
silberlingi, Picrodus, 224.
siluricola, Arhythmorhynchus, 153.
Simaruba miocenica, 338, 340.
similaris, Rileya, 482.
similis, Haruspex, 196.
similis, Perilampus, 371, 376, 406.
simplex, Hyopsodus, 242.
Simplicella glacialis, 443.
simplicidens, Plesippus, 282, 288, 289, 292, 293, 316, 318–320.
simplicidens, *Stilpnodon*, 223, 229.
Simpson, George Gaylord, on new Paleocene mammals from Fort Union of Montana, 221.
simpsoni, Parectypodus, 227.
sinclairi, Ptilodus, 223, 225, 226.
sinensis, Marphysa, 261, 266.
Nephthys, 261, 276.
singularis, Atrophopoda, 17, 18, 38.
Paradidyma, 10, 15, 19, 21, 31, 32, 36–38, 41.
sinuosa, Basilemys, 180, 181.
Siphonacme, 364.
lyttoni, 364.
Siphonophora, 361, 362, 364.
limitare, 362, 363 (fig.).
Sisymbrium, 407.
Skinker, Mary Scott, on new tapeworms from carnivores and redescription of Taenia laticollis Rudolphi, 211.
smithi, Oxyurostylis, 423, 437, 438.
Smulyan, Marcus Thomas, on chalcid flies of genus Perilampus, 369.
solitarium, Nymphon, 414, 417 (fig.).
Sophora, 350.
marana, 338, 340, 346 (fig.), 350.
salvadorana, 338, 340.
Southwellina, 153.
Spalacopsis grandis, 201.
ornatipennis, 200.
Spanipalpus, 46.
aldrichi, 52.
australis, 57.
miscelli, 49.
Spanoxyodon, 236.
latrunculus, 224, 236.
spatula, Pliauchenia, 288.
speciosa, Tentacularia, 130.
speciosum, Rhynchobothrium, 130.
spectabilis, Cassia, 349.
Sphaeralcia angustifolia, 407.
sphaerocarpa, Anona, 347.
sphaerocarpoides, Anona, 338, 340, 346 (fig.), 347.
spiculigerum, Contracaecum, 143.
spiracornuta, Tentacularia, 130.
Spirastrella, 461.
cunctatrix, 461.
Sponges, shallow-water, from near Panama Canal, 441.
Spongia, 455.
barbara, 455.
campana, 457.
cinerea, 444, 445.
lignea, 455.
panicea, 449.
Squatinidae, 185 (fig.).
Stegomastodon successor, 288.
Steigstrum, 331.
Stenosmilia, 76.
protelaria, 76.
Stephanocoenia sp., 104.

stephensoni, Combretum, 338, 340, 355, 359 (fig.).
sternbergii, Pentaceratops, 160, 163, 185.
Stiboriopsis jamaicaensis, 72.
Stilpnodon, 229.
simplicidens, 223, 229.
Stlengis *distoechus*, 325, 327, 328–330 (fig.).
misakia, 327, 330.
osensis, 325–328.
revision of genus, 325, 326.
stokesi, Dichocoenia, 75.
strenuana, Epiblema, 405.
stricklandi, Cuphocera, 63.
Trichophora, 63.
Strongylophora, 459.
durissima, 459.
santa, 459.
stygicus, Perilampus, 371, 376, 404, 411.
stylifer, Hippocampus, 499, 561, 566, 567.
Stylinodontidae, 223.
Stylocoenia, 105.
Stylophora, 93, 94.
cambridgensis, 73, 95, 110.
contorta, 73, 94.
sp. *a* and *b*, 73, 93.
Styracaceae, fossil Venezuelan, 356.
Styrax, 356.
lanceolata, 340, 356.
subcarinatus, Perilampus, 370, 373, 386.
Suberites sulcatus, 450.
subtriangularis, Thalysias, 458.
subtrigonus, Tricentes, 237.
successor, Stegomastodon, 288.
sulcatus, Pseudosuberites, 442, 449.
Suberites, 450.
superior, Tetraclaenodon, 224, 239.
Sweetia, 349.
Sylvilagus, 114–117, 120.
nuttalli grangeri, 111, 114–116.
symbolicus, Tetraclaenodon, 224, 239, 240.
Sympherobius angustus, 395, 411.
Synastrea, 87.
agaricites, 88.
adkinsi, 73, 83, 87, 109.
Synbothrium, 130, 132.
filicolle, 130.
fragile, 130.
hemuloni, 130.
malleum, 133.
Syndesmobothrium, 131.
filicolle, 131.
fragile, 130.
Syngnathoides biaculeatus, 515.
Syngnathus erectus, 516, 517.
foliatus, 515, 516.
hippocampus, 513–520, 525, 530, 543, 570.
longirostris, 522.
tetragonus, 515.
synodi, Lecithochirium, 128.
Taboga, 452.
taboga, 452, 453 (fig.).
Tachina macrocera, 52.
Taenia crassicollis, 217, 218.
laticollis, redescription of, 211, 216.
lyncis, 211, 212.
macrocystis, 214, 217, 219.
taeniaeformis, 217, 218.
taxidiensis, 211, 215.
taeniaeformis, Taenia, 217, 218.
Taeniodonta, Fort Union, 223.
Tanaorhamphus, 151.
Tanupolama, 285, 288.
Tapeworms, new species from carnivores, 211.
Tapirira, 352.
lanceolata, 340, 352, 353 (fig.), 360.
trinitiana, 340, 352, 353 (fig.).
tardus, Parectypodus, 225.
Taxidea taxus taxus, 215.
taxidiensis, Taenia, 211, 215.
Taxocrinus, 4.
briareus, 4, 5.
taxus, Taxidea taxus, 215.
Tedania, 459.
ignis, 459.
tener, Leptacodon, 228.
Tennessee, millipeds from, 361.
Tentacularia *lepida*, 125, 129.
speciosa, 130.
spiracornuta, 130.
tenuicornis, Rhadinorhynchus, 123, 125, 154.
tenuipes, Adelometra, 247.
Caryometra, 247.
tenuis, Diastylopsis, 436, 437 (fig.).
Didymictis, 224, 238.
Terebellidae, 278.
Tertiary, new Foraminifera of genera Operculina and Operculinoides from North America and West Indies, 487.
Venezuelan plants from, 335.
Testudinata, fossil, from Idaho, 285.
Tethya, 451, 452, 462.
aurantia, 462.
diploderma, 451.
Tethyidae, 452.
Tetraclaenodon, 239-241.
pliciferus, 239.
superior, 224, 239.
symbolicus, 224, 239, 240.
tetragonus, Syngnathus, 515.
tetraphylla, Inga, 349.
tetraquetrum, Pleonotoma, 358.
Tetrarhynchidae, 129.
Tetrarhynchus erinaceus, 132.
platycephalus, 130, 131, 133.
Tetrastichus, 485.
gerstaeckeriae, 485.
malacosomae, 485, 486.
texanum, Glyptotherium, 288.
texanus, Platygonus, 288.
Texas, millipeds from, 361.
parasites of fishes from Galveston Bay, 123.
texensis, Apocynophyllum, 341.
Thalysias ignis, 459.
subtriangularis, 458.

Thamnasteria, 88.
Thamnophis sp., 285.
Thelepus, 278.
japonicus, 278.
Thelvetia, 357.
Thescelus, 170, 172, 175, 187.
fluviatilis, 177.
hemispherica, 174, 175 (fig.), 176 (fig.), 177, 186.
insiliens, 174–177.
rapiens, 174, 175, 177.
Thomomys gidleyi, 285, 288.
Thryptacodon, 233, 234.
tobleri, Lepidocyclina, 490.
tongi, Hernandia, 340, 353 (fig.), 354.
torosa, Cuphocera, 46, 48, 67.
Torpedo, 134.
townsendi, Atrophopoda, 10, 12.
Ceratomyiella, 11, 12.
Oedemapeza, 12.
Paradidyma, 12.
townsendii, Lepus, 113, 120.
Lepus townsendii, 112.
Toxadocia, 445.
proxima, 442, 445.
transversale, Rhipidocotyle, 125, 126, 128, 141.
Treadwell, Aaron Louis, on polychaetous annelids from Amoy, China, 261.
Trechmann, Charles Taylor, his collection of fossil corals, 74, 93, 94, 103.
trechmanni, Dichocoenia, 72, 75, 109.
Goniopora, 73, 91, 109.
Trematoda, 126.
new genus and species from little brown bat and key to Pleurogenetinae, 321.
triacis, Orthonopias, 326.
triangularis, Perilampus, 411.
Tricentes, 233, 236.
latidens, 224, 236.
subtrigonus, 237.
trichiuri, Porrocaecum, 125, 145.
Trichiurus lepturus, 125, 128, 145.
Trichophora, 70.
fucata, 65.
miscelli, 46, 49.
stricklandi, 63.
tricuspe, Contracaecum, 143.
tridentatum, Nymphon, 418.
Trigonia, 351.
varians, 339, 340, 351, 353 (fig.).
Trigoniaceae, fossil Venezuelan, 351.
triloculatum, Phoreiobothrium, 134.
trimaculatus, Hippocampus, 517, 518.
trinitiana, Tapirira, 340, 352, 353 (fig.).
trinitatis, Lepidocyclina, 490.
Trionychidae, 165, 182, 186.
triphylla, Bignonia, 358.
Trocharaea, 86.
Trochocyathus, 74, 104, 105.
matleyi, 72, 74, 109.
sp., 103, 105.
woolmani, 74.
Trochoplegma, 86.
Trochoseris, 78, 81, 88, 98.
catadupensis, 72, 78.
sp., 73, 98.
Trochosmilia hilli, 78, 79.
tropicus, Cordillerion, 288.
trovessartianus, Ptilodus, 225.
Trypespongia, 456.
columbia, 456.
tuberculata, Nanilla, 196.
Operculina, 487, 488, 495.
tuberosa, Dichocoenia, 75.
tubicen, Parasaurolophus, 161, 185.
tunicatae, Nereis (Ceratonereis), 476, 477 (fig.).
Turbinoseris cantabrigiensis, 96.
jamaicaensis, 97.
turritum, Nymphon, 414, 416, 417 (fig.).
Turtles, fossil, from Kirtland formation of New Mexico, 159.
tuxpanicus, Operculinoides, 488, 494, 495.
tylodermae, Neocatolaccus, 484.
Tylosurus marinus, 126.
uinta, Lynx rufus, 214.
ulkei, Monellema, 485.
undatum, Chisternon, 172.
undulatus, Micropogon, 125, 131, 154.
unicornis, Cyclaspis, 427.
Unionidae, 127.
uniplicata, Hemilamprops, 431.
Ursus, 286.
vagus, Alilepus, 119 (fig.), 121, 285, 288.
valida, Eulimneria, 382, 411.
validinervis, Didyma, 17, 37.
Paradidyma, 20, 23, 25, 29, 33–35, 37, 39, 40, 42, 43.
vanduzeei, Leptostylus, 204.
Vanessa cardui, 252.
iortlandia, 252.
itea, 252.
portlandia, 252.
variabilis, Hircinia, 457.
varians, Cyclaspis, 427.
Trigonia, 339, 340, 351, 353 (fig.).
variegatus, Cyprinodon, 123–125, 140–142, 150, 152.
variolosa, Basilemys, 180, 181.
Vaughan, Thomas Wayland, and Cole, W. Storrs, on new Tertiary Foraminifera from North America and West Indies, 487.
Vaughanoseris, 80.
catadupensis, 72, 80, 81, 109, 110.
vecordensis, Claenodon, 223, 232.
vegetus, Aspideretes, 183.
Velates schmiedeliana, 95, 101.
Venezuela, Tertiary plants from, 335.
venezuelana, Burserites, 338, 340.
venezuelanus, Chrysobalanus, 338, 340, 342 (fig.), 347.
venezuelensis, Antholithus, 338, 340.
Diplaraea, 84.
Leguminosites, 338, 340.
venosa, Operculinella, 489.
verecundus, *Atactorhynchus*, 123, 125, 151.

vetus, Hypolagus, 112 (fig.), 114, 116, 117, 119-121, 285, 288.
vexillosa, Nereis, 469–471.
vicksburgensis, Operculinoides, 487, 490, 495.
villosoides, Coussapoa, 340.
villosus, Hippocampus, 529, 582, 583.
violacea, Chalcis, 371.
Diplolepis, 371.
virens, Nereis (Neanthes), 479.
virginianus, Odocoileus, 214.
viridis, Orasema, 372.
vomitor, Agamonema, 125, 150.
vorax, Aspideretes, 183, 184 (fig.), 185, 186.
walkeri, Parasaurolophus, 162, 163.
walli, Astreopora, 73, 102, 110.
Washington, pycnogonids from Puget Sound, 413.
Wells, John West, on fossil corals from West Indies, 71.
West Indies, new cerambycid beetles from, 189.
new Foraminifera from Tertiary of, 487.
fossil corals from, 71.
wickhami, Elaphidion, 192.
Eupogonius, 199.
wilcoxensis, Ficus, 343.
willcoxi, Nummulites, 489.
Willughby, Francis, his account of seahorses, 519.
wolfi, Meniscium, 340, 342 (fig.), 343.
woolmani, Trochocyathus, 74.
Worms, parasites of Galveston Bay fishes, 123.
polychaetous annelids from Amoy, China, 261.
new polychaetous annelids (Nereidae) from California, 467.
new tapeworms from carnivores, 211.
new trematode from little brown bat and key to Pleurogenetinae, 321.
xanthurus, Leiostomus, 125, 148, 154.
Zamia, 344.
sp., 338, 340, 344.
Zenillia (Exorista) sp., 393, 403, 411.
Zenometrinae, 247, 248.
zeteki, Laxosuberites, 450, 451 (fig.), 455.
Zetekispongia, 446.
zonca, 446, 447 (fig.).
Zimmer, Carl, on California Crustacea of order Cumacea, 423.
zonca, *Zetekispongia*, 446, 447 (fig.).
zosterae, Hippocampus, 502, 506, 512, 525, 529, 585, 588, 589.
zuliana, Bignonia, 338, 340, 358, 359 (fig.).
zuliana, Cassia, 340, 346 (fig.), 350.
zyggompha, Roosa, 462.

U. S. GOVERNMENT PRINTING OFFICE: 1956